



Planta Medica

Journal of Medicinal Plant and Natural Product Research

11

Volume 78
August 2012
Page 1027-1310

8th Joint Meeting of AFERP, ASP, GA, PSE & SIF

28th July–1st August 2012, New York City, USA

Abstracts

Chairmen:

Guy T. Carter

Edward J. Kennelly

Issue Editor:

Mark O'Neil-Johnson

PF56

Plant species from the Peruvian Amazon rainforest (Peru) and their antimicrobial activity
 Roumy V¹, Gutierrez-Choquevilca AL², Lopez Mesia JP³, Ruiz L³, Ruiz J³, Abedini A¹, Hennebelle T¹, Neut C⁴
¹Laboratoire de Pharmacognosie, EA 4481, Université de Lille, 59006 Lille, France; ²Laboratoire EREA, LESC, UMR 7186 CNRS Villejuif/Université Paris Ouest-Nanterre La Défense, France; ³Universidad Nacional de la Amazonia Peruana (UNAP), Iquitos, Perú; ⁴Laboratoire de Bactériologie, U995, Université de Lille, 59006 Lille, France

The plant species reported here are traditionally used by Indigenous and Mestizo populations from the Iquitenian surroundings (Peruvian Amazon) for microbial infections. Inhabitants of various ethnic origins were interviewed and selected plants extracts were evaluated for their antimicrobial properties against 36 sensitive and multi-resistant bacteria or fungi. Of the 39 plants analyzed (50 methanolic extracts), 9 species showed MIC ≤ 0.3 mg/ml for one or several microorganisms and only 6 extracts were inactive. This study supports the traditional use of these plants. It may help to discover new chemical classes of antibiotics that could serve as selective agents against multi-resistant bacteria.

PF57

***Cistus ladanifer* as a source of phenolic compounds with antifungal activity**

Barros L^{1,2}, Dueñas M², Alves CT³, Silva S³, Henriques M³, Santos-Buelga C², Ferreira JCFR¹
¹CIMO-ESA Polytechnic Institute of Bragança, Portugal; ²GIP Faculty of Pharmacy, University of Salamanca, Spain; ³IBB Centre of Biological Engineering, University of Minho, Braga, Portugal

A screening of the antifungal potential of phenolic extract of *Cistus ladanifer* from Northeast Portugal, against *Candida* species was performed. The extract was characterized by HPLC-DAD-ESI/MS. Phenolic acids and derivatives, ellagic acid derivatives and flavonoids, such as catechins, flavonols and flavones, were found in the sample. The most abundant group was ellagic acid derivatives in which punicalagin gallate, a derivative of punicalagin attached to gallic acid, was found in highest amount. These compounds could be related to the strong inhibition of *C. albicans*, *C. glabrata* and *C. parapsilosis* growth. Moreover, the best antifungal activity was against *C. glabrata*, where the studied extract was able to cause at least 3 Log of reduction at concentrations below 50 µg/mL and a total growth inhibition at concentrations above 625 µg/mL.

PF58

Effects of *Ocimum sanctum* Linn (OS) leaf extract on stress, memory and attention in healthy humans

Suneetha S, Talwar A, Mahapatra SC, Sharma R
 Dept of Physiology, All India Institute of Medical Sciences, New Delhi, India 110029

Anti-stress effects and memory enhancing effects of OS have been documented in animal models but no human studies available. Double blinded RCT on healthy human adults. 300 mg capsules of ethanolic leaf extract of OS or placebo were administered to 30 volunteers for 30 days. Recordings were taken on day 0, 15 and 30. Parameters assessed: a) STAI questionnaire b) Sternberg memory task c) Stroop task d) Heart rate (HR) & e) GSR. Results: showed significant improvement in reaction time $p=0.043$ in Sternberg task, and improved % of correct responses in facilitation task of Stroop ($p=0.01$). STAI, HR, GSR and P300 latency showed no significant change. P300 amplitude showed a significant increase ($p=0.02$). All changes were significant only at day 30. OS possesses memory enhancing effects at dose of 300 mg/day od by 30 days. There is also significant improvement in attention as assessed by P300 amplitude. However no significant reductions in Stress parameters were seen at this dose/duration of OS. Further trials have to be conducted in more subjects and for longer duration for the effects to be validated

PF59

Phytochemical investigation of *Rauvolfia nukuhivensis*, a Marquesan traditional medicinal plant

Martin N¹, Thomas O², Prado S³, Paetz C⁴, Lecellier G¹, Raharivelomanana P¹
¹BIOTEM, University of French Polynesia, BP 6570 Faaa, 98702 FAA'A, Tahiti, French Polynesia; ²ICN UMR 7272, University of Nice Sophia-Antipolis, Parc Valrose 06108 Nice Cedex 2, France; ³MNHN UMR CMAM 7245, 63 rue Buffon, 75230 Paris, France; ⁴MPI for Chemical Ecology, Hans-Knöll-Strasse 8, 07745 Jena, Germany



Rauvolfia nukuhivensis (Apocynaceae) is an endemic species of the Marquesas archipelago where it is used as a traditional gynecological anti-septic. Over-exploited because of the frequent use of the bark (macerate), the plant is now classified as an endangered species ("Critical Rare" UICN status). Data regarding pharmacological principles and their chemical identity were not available until now. The phytochemical investigation of the main constituents of this popular medicinal plant resulted in the isolation and identification of several alkaloids belonging to the sandwicine and ajmaline type, among them also formerly unknown derivatives. In order to test the efficacy against human pathogens (*Staphylococcus aureus*, *Escherichia coli* and *Candida albicans*) bioassays were carried out, showing moderate antifungal activities of some compounds.

PF60

Analysis of leaf epidermal characters of medicinal and poisonous Brazilian menispermaceae

Porto NM, Araújo ND, Basílio IJLD, de Fátima Agra M
 PPGPNSB, Universidade Federal da Paraíba, CP 5009, 58015-970, João Pessoa, Paraíba, Brazil

We carried out a multivariate analysis of leaf anatomy of five genera and nine species of Menispermaceae known to be toxic or used as medicinal in Brazil. The dendrogram obtained by Ward's method showed two groups (A and B), with cophenetic correlation coefficient of 0.9178. The group A with six species and two subgroups: the subgroup A1 brought together four species of *Cissampelos* with anomocytic stomata (Fig 1A), without inclusions of calcium oxalate (Fig.1E) and sclereids; the subgroup A2 with *Cissampelos pareira*, *Odontocarya* and *Disciphania*. The group B consists of *Chondrodendron platyphyllum* with anticlinal walls cells straight (Fig 1G) and *Hyperbaena domingensis* with cyclocytic stomata (Fig.1B), bractiform cells in the spongy parenchyma (Fig 1C), and secretory canals (Fig 1D). *Cissampelos sympodialis* and *Chondrodendron platyphyllum* were the species with lower similarity in the dendrogram. Financial support: CNPq.



A. anomocytic stomata; B. cyclocytic stomata; C. bractiform cells in the spongy parenchyma; D. secretory canals; E. inclusions of calcium oxalate; F. Papillae; G. Anticlinal walls cells straight.